

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 45/06, 31/44, 31/445, 9/20, 9/26, 9/48</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/25065</b>
			<b>(43) International Publication Date:</b> 17 July 1997 (17.07.97)
<b>(21) International Application Number:</b> PCT/SE96/01736			<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
<b>(22) International Filing Date:</b> 20 December 1996 (20.12.96)			
<b>(30) Priority Data:</b> 9600072-4 8 January 1996 (08.01.96) SE			
<b>(71) Applicant (for all designated States except US):</b> ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE).			
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> DEPUI, Helene [FR/SE]; Wrangelsgatan 7B, S-416 62 Göteborg (SE). HALLGREN, Agneta [SE/SE]; Hökegårdsgatan 2C, S-431 38 Mölndal (SE).			
<b>(74) Agent:</b> ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).			<b>Published</b> With international search report.

**(54) Title:** ORAL PHARMACEUTICAL DOSAGE FORMS COMPRISING A PROTON PUMP INHIBITOR AND A PROKINETIC AGENT

**(57) Abstract**

An oral pharmaceutical dosage form comprising a proton pump inhibitor and one or more prokinetic agents in a fixed formulation, wherein the proton pump inhibitor is protected by an enteric coating layer. The fixed formulation is in the form of multilayered tablets, capsules or multiple unit tableted dosage forms. The multiple unit dosage forms are most preferred. The new fixed formulation is especially useful in the treatment of disorders associated with gastro oesophageal reflux diseases.



BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## ORAL PHARMACEUTICAL DOSAGE FORMS COMPRISING A PROTON PUMP INHIBITOR AND A PROKINETIC AGENT

### Field of the invention

5 The present invention is related to new oral pharmaceutical preparations especially for use in the prevention and treatment of disorders associated with gastro oesophageal reflux. The present preparations comprise a gastric acid suppressing agent, such as a proton pump inhibitor, in combination with one or more prokinetic agents in a new fixed unit dosage form, especially a tablet. Furthermore, the present invention refers to a method for the  
10 manufacture of such preparations and the use of such preparations in medicine, especially in the treatment of gastro oesophageal reflux diseases and other gastrointestinal disorders.

### Background of the invention

15 Gastro oesophageal reflux disease (GORD) is among the most common disorders seen by gastroenterologists and general practitioners. The wide diversity of symptoms and disease severity produced by acid reflux has led to the need for more individualized treatment strategies. Therapeutic agents effective in the treatment of GORD include gastric acid suppressing agents, such as H<sub>2</sub> receptor antagonists, proton pump inhibitors, other agents  
20 of interest are antacids/alginates and prokinetic agents. These agents can be distinguished by their mechanisms of action, safety profile, pharmacokinetics and indications.

Antacids and alginates are still widely used. They have a short duration of action but are seen as inexpensive and safe. They do not provide a layterm symptom resolution of GORD.

25

H<sub>2</sub> receptor antagonists are widely prescribed for GORD. Their higher cost has been compensated by the clinical results obtained both in terms of symptom relief and healing. These advantages have been related to their mode of action, which offered more potent and longer duration of effect on gastric acidity.

Proton pump inhibitors, such as omeprazole, are rapidly taking share from H<sub>2</sub> receptor antagonists, particularly in reflux oesophagitis. Omeprazole is known to offer significant gain over H<sub>2</sub> receptor antagonists in terms of symptom resolution, healing and prevention of relapse for reflux oesophagitis.

Prokinetic agents of the first generation, e.g. bethanecol, stimulates cholinergic receptors, and of the second generation, e.g. domperidone and metoclopramide, blocks effects of endogenous dopamine in the gut. The results of double-blind placebo controlled trials in GORD patients have been conflicting. The action of the third generation of prokinetic agents, such as substituted benzamides, e.g. cisapride and mosapride derives primarily, but not exclusively, from facilitating acetylcholine release from neurones of the myenteric plexus via stimulation of 5-HT<sub>4</sub> receptors. The efficacy of orally administered benzamides, such as cisapride, in patients with GORD and reflux oesophagitis has been studied and a superior effect in alleviating gastro-oesophageal symptoms and healing low grade oesophagitis (non circumferential erosion) has been shown in most studies.

Patients with severe symptoms, severe mucosal damage or both are almost always treated with proton pump inhibitors for profound and long-term control of gastric acid secretion. Patients with mild symptoms and limited mucosal damage respond best to H<sub>2</sub>-receptor antagonist, prokinetic agents or proton pump inhibitors.

A combination therapy of a prokinetic agent and a gastric acid lowering compound is rational and was shown more effective than mono therapy apart from full dose of proton pump inhibitors. Administration of cisapride and ranitidine was shown to further lower the exposure of the oesophagus to acid(s) (Inauen W et al. Gut 1993; 34: 1025 - 1031). Such a therapy was also shown to improve healing rates (de Boer WA et al. Aliment Pharmacol Ther 1994; 8: 147 - 157). WO 95/01803 describes a pharmaceutical composition of famitidine, cisapride and optionally simethicone in the treatment of gastrointestinal distress.

Maintenance therapy is often necessary to prevent recurrent symptoms and oesophagitis. Recently a combination therapy combining an acid-suppressing medication with a prokinetic (cisapride) was shown also very effective. Further, Vyneri et al (N. Engl. J Med 1995; 333: 1106 - 1110) found that omeprazole alone or in combination with cisapride was more  
5 effective than ranitidine alone or cisapride alone and that omeprazole combined with cisapride was more effective than ranitidine plus cisapride. Such combination therapies might be considered for patients whose predominant symptom is regurgitation; those whose symptoms occur mainly at night; those with respiratory problems such as posterior laryngitis, asthma, chronic bronchitis, or recurrent aspiration; those with cough and  
10 hoarseness related to reflux disease.

A combination therapy comprising an acid suppressing agent and a prokinetic agent is attractive, rational and effective. An acid suppressing agent plus a prokinetic agent could be an alternative to each of them separately in case of failure. However, because of the large  
15 number of therapeutical tablets/pills that must be taken each day in such a therapy, the compliance of such a treatment may be a problem. It is well known that patient compliance is a main factor in receiving good results in medical treatments. Administration of two, three or even more different tablets to the patient is not convenient or satisfactory to achieve the most optimal results. The present invention now provides new oral dosage forms  
20 comprising two or more different active substances combined in one fixed unit dosage form, preferably a tablet.

It is well known that some of the gastric acid suppressing agents, such as proton pump inhibitors are susceptible to degradation/transformation in acid reacting and neutral media.  
25 In respect of the stability properties, it is obvious that the one of the active substances being an acid susceptible proton pump inhibitor must be protected from contact with acidic gastric juice by an enteric coating layer. There are different enteric coating layered preparations of proton pump inhibitors described in the prior art, see for example US-A 4,786,505 (AB Hässle) describing a preparation comprising omeprazole.

There are problems to produce a fixed unit dosage form comprising a rather high amount of active substance. Different active substances with differing physical properties in the same preparation give further problems. Preparation of a multiple unit tableted dosage form arises specific problems when enteric coating layered pellets containing acid susceptible proton pump inhibitors as active substance are compressed into tablets. If the enteric coating layer does not withstand the compression of the pellets into a tablet the susceptible active substance will be destroyed by penetrating acidic gastric juice, i.e. the acid resistance of the enteric coating layer of the pellets will not be sufficient in the tablet after compression.

#### 10 Summary of the invention

The present invention provides oral, fixed unit dosage forms, i.e. a multiple unit tableted dosage forms, multilayered tablets or a capsule filled with more than one pharmaceutically active compound. The active compounds present in the dosage form are preferably an acid susceptible proton pump inhibitor which is protected by an enteric coating layer, and one or more prokinetic agents. These new dosage forms will simplify the regimen and improve the patient compliance.

#### Brief description of the Figures

20

Fig. 1 illustrates a cross-section of a multiple unit tableted dosage form comprising an acid susceptible proton pump inhibitor in the form of enteric coating layered pellets (1) in admixture with a prokinetic agent and pharmaceutically acceptable excipients (2). The tablet is covered by a filmcoating layer, i.e. tablet coat (7).

25

Fig. 2 illustrates a cross-section of a tablet with two separate layers, one of which comprising enteric coating layered pellets (1) in admixture with excipients (3) and the other layer comprising the prokinetic agent in admixture with pharmaceutically acceptable excipients (2). The tablet is covered by a filmcoating layer (7).

30

Fig. 3 illustrates a cross-section of an enteric coating layered tablet comprising a proton pump inhibitor in admixture with pharmaceutically acceptable excipients in the tablet core (5) surrounded by an enteric coating layer (8) and thereupon a layer of the prokinetic agent(s) in admixture with pharmaceutically acceptable excipients (6). The tablet is covered by a filmcoating layer (7).

Fig. 4 illustrates a cross-section of a multiple unit tableted dosage form comprising an acid susceptible proton pump inhibitor in the form of enteric coating layered pellets (1) in admixture with excipients (3) and on the multiple unit tableted dosage form a layer comprising the prokinetic agent(s) in admixture with pharmaceutically acceptable excipients (6). The tablet is covered by a filmcoating layer (7).

#### Detailed description of the invention

One object of the invention is to provide an oral, multiple unit tableted dosage form comprising an acid susceptible proton pump inhibitor in the form of individually enteric coating layered units together with one or more prokinetic agents in the form of a powder or granules compressed into a tablet. The enteric coating layer(s) covering the individual units of the proton pump inhibitor has properties such that the compression of the units into a tablet does not significantly affect the acid resistance of the individually enteric coating layered units. Furthermore, the multiple unit tableted dosage form provides a good stability of the active substances during long-term storage.

The new fixed dosage form is preferably in the form of a multiple unit tableted dosage form comprising enteric coating layered units of the one of the active substance which is acid susceptible and granules of the other active substance, i.e. prepared prokinetic granules as shown in Fig. 1.

The proton pump inhibitor, in the form of enteric coating layered units, may also be mixed with pharmaceutically acceptable excipients and compressed into a tablet which is then filmcoated with an aqueous suspension containing the prokinetic substance, see Fig. 4.

5 Another object of the invention is to provide a tablet preparation comprising a proton pump inhibitor in admixture with tablet excipients in a tablet core and a separate layer surrounding the tablet core, which layer comprises one or more prokinetic agent(s) presscoated onto the tablet core. The tablet core is enteric coating layered before the surrounding layer of prokinetic agents is applied. Optionally a separating layer also is applied on the tablet before  
10 the enteric coating layer, see Fig. 3.

Alternatively, the prepared tablet is sectioned in separate layers, each one comprising different active substances. Preferably one layer comprises the proton pump inhibitor in the form of enteric coating layered pellets in admixture with pharmaceutically acceptable  
15 excipients and another layer(s) comprises(-e) the prokinetic agent(s) in admixture with pharmaceutically acceptable excipients, respectively, see Fig. 2.

A further object of the invention is to provide a multiple unit tableted dosage form, which is divisible and easy to handle. Such a multiple unit tableted dosage form may be dispersed in  
20 an aqueous liquid and can be given to patients with swallowing disorders and in pediatrics. Such a suspension of dispersed units/pellets of appropriate size can be used for oral administration and also for feeding through a naso-gastric tube.


Furthermore, the present invention provides a capsule preparation comprising the proton  
25 pump inhibitor in the form of enteric coating layered pellets mixed with one or more prokinetic agents in the form of prepared granules or pellets. The new fixed unit dosage forms comprise as active substances one gastric acid suppressing agent, such as an acid susceptible proton pump inhibitor and one or more prokinetic agents. The different therapeutically active components used in the dosage forms are defined below.



The prokinetic part of the formulation may be formulated in the form of instant release, sustained release or extended release formulations. Alternatively, all the components of the formulation may be formulated in an effervescent formulation.

# 5 Active substances


The gastric acid suppressing agent is preferably an acid susceptible proton pump inhibitor. Such proton pump inhibitors are for example compounds of the general formula I




10

wherein

Het<sub>1</sub> is




or




15


Het<sub>2</sub> is



or




or




20

X =



or



wherein

- 5 N in the benzimidazole moiety means that one of the carbon atoms substituted by R<sub>6</sub>-R<sub>9</sub> optionally may be exchanged for a nitrogen atom without any substituents;

R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are the same or different and selected from hydrogen, alkyl, alkoxy optionally substituted by fluorine, alkylthio, alkoxyalkoxy, dialkylamino, piperidino,  
10 morpholino, halogen, phenyl and phenylalkoxy;

R<sub>4</sub> and R<sub>5</sub> are the same or different and selected from hydrogen, alkyl and aralkyl;

R<sub>6</sub>' is hydrogen, halogen, trifluoromethyl, alkyl and alkoxy;


15 R<sub>6</sub>-R<sub>9</sub> are the same or different and selected from hydrogen, alkyl, alkoxy, halogen, haloalkoxy, alkylcarbonyl, alkoxycarbonyl, oxazolyl, trifluoroalkyl, or adjacent groups R<sub>6</sub>-R<sub>9</sub> form ring structures which may be further substituted;

R<sub>10</sub> is hydrogen or forms an alkylene chain together with R<sub>3</sub> and



20


R<sub>11</sub> and R<sub>12</sub> are the same or different and selected from hydrogen, halogen or alkyl, alkyl groups, alkoxy groups and moities thereof, they may be branched or straight C<sub>1</sub> - C<sub>9</sub> - chains or comprise cyclic alkyl groups, such as cycloalkylalkyl.

25 Examples of proton pump inhibitors according to formula I are




Omeprazole






Lansoprazole




Pantoprazole





5




Pariprazole



Leminoprazole





5 The proton pump inhibitors used in the dosage forms of the invention may be used in neutral form or in the form of an alkaline salt, such as for instance the  $Mg^{2+}$ ,  $Ca^{2+}$ ,  $Na^+$ ,  $K^+$  or  $Li^+$  salts, preferably the  $Mg^{2+}$  salts. Further where applicable, the compounds listed above may be used in racemic form or in the form of a substantially pure enantiomer thereof, or alkaline salts of the single enantiomers.

10 Suitable proton pump inhibitors are for example disclosed in EP-A1-0005129, EP-A1-174 726, EP-A1-166 287, GB 2 163 747 and WO90/06925, WO91/19711, WO91/19712, and further especially suitable compounds are described in WO95/01977 and WO94/27988.

15 The gastric acid suppressing agent is preferably an acid susceptible proton pump inhibitor but other gastric acid suppressing agents such as the  $H_2$  receptor antagonists: ranitidine, cimetidine or famotidine, may be used together with a prokinetic agent in the pharmaceutical compositions according to the present invention.

20 A wide variety of prokinetic compounds may be used in combination with a suitable proton pump inhibitor in the fixed unit dosage form according to the present invention. Such prokinetic agents include for example cisapride, mosapride, metoclopramide, and domperidone. The active prokinetic agents could be in standard forms or used as salts, hydrates, esters etc. A combination of two or more of the above described drugs may be

used. A preferable prokinetic agent for the new fixed dosage form is mosapride or cisapride. Such suitable prokinetic agents are described in EP 0 243 959 and EP 0 076 530.

5 The preferred multiple unit tableted dosage form comprising a proton pump inhibitor in the form of a racemat, an alkaline salt or one of its single enantiomers in combination with a prokinetic compound, is characterized in the following way. Individually enteric coating layered units (small beads, granules or pellets) containing the proton pump inhibitor and optionally alkaline reacting substances, are mixed with the prokinetic compound and conventionally tablet excipients. The prokinetic compound and tablet excipients may be dry  
10 mixed or wet-mixed into granules. The mixture of enteric coating layered units, prokinetic agent(s) and optionally excipients are compressed into the multiple unit tableted dosage forms. With the expression "individual units" is meant small beads, granules or pellets, in the following referred to as pellets of the proton pump inhibitor.

15 The compaction process (compression) for formulating the multiple unit tableted dosage form must not significantly affect the acid resistance of the enteric coating layered pellets. In other words the mechanical properties, such as the flexibility and hardness as well as the thickness of the enteric coating layer(s), must secure that the requirements on enteric coated articles in the United States Pharmacopeia are accomplished in that the acid resistance does  
20 not decrease more than 10% during the compression of the pellets into tablets.

The acid resistance is defined as the amount of proton pump inhibitor in the tablets or pellets after being exposed to simulated gastric fluid USP, or to 0.1 M HCl (aq) relative to that of unexposed tablets and pellets, respectively. The test is accomplished in the following  
25 way. Individual tablets or pellets are exposed to stimulated gastric fluid of a temperature of 37°C. The tablets disintegrate rapidly and release the enteric coating layered pellets to the medium. After two hours the enteric coating layered pellets are removed and analyzed for content of the proton pump inhibitor using High Performance Liquid Chromatography (HPLC).

Further specific components used in the fixed unit dosage forms of the present invention are defined below.

Core material - for enteric coating layered pellets comprising a proton pump inhibitor

- 5
- The core material for the individually enteric coating layered pellets can be constituted according to different principles. Seeds layered with the proton pump inhibitor, optionally mixed with alkaline substances, can be used as the core material for the further processing.
- 10
- The seeds which are to be layered with the proton pump inhibitor can be water insoluble seeds comprising different oxides, celluloses, organic polymers and other materials, alone or in mixtures or water-soluble seeds comprising different inorganic salts, sugars, non-pareils and other materials, alone or in mixtures. Further, the seeds may comprise the proton pump inhibitor in the form of crystals, agglomerates, compacts etc. The size of the seeds is not
- 15
- essential for the present invention but may vary between approximately 0.1 and 2 mm. The seeds layered with the proton pump inhibitor are produced either by powder or solution/suspension layering using for instance granulation or spray coating layering equipment.
- 20
- Before the seeds are layered, the proton pump inhibitor may be mixed with further components. Such components can be binders, surfactants fillers, disintegrating agents, alkaline additives or other and/or pharmaceutically acceptable ingredients alone or in mixtures. The binders are for example polymers such as hydroxypropyl methylcellulose (HPMC), hydroxypropylcellulose (HPC), carboxymethylcellulose sodium, polyvinyl
- 25
- pyrrolidone (PVP), or sugars, starches or other pharmaceutically acceptable substances with cohesive properties. Suitable surfactants are found in the groups of pharmaceutically acceptable non-ionic or ionic surfactants such as for instance sodium lauryl sulfate.
- 30
- Alternatively, the proton pump inhibitor optionally mixed with alkaline substances and further mixed with suitable constituents can be formulated into a core material. Said core



material may be produced by extrusion/spheronization, balling or compression utilizing conventional process equipment. The size of the formulated core material is approximately between 0.1 and 4 mm and preferably between 0.1 and 2 mm. The manufactured core material can further be layered with additional ingredients comprising the proton pump inhibitor and/or be used for further processing.

The proton pump inhibitor is mixed with pharmaceutical constituents to obtain preferred handling and processing properties and a suitable concentration of the substance in the final mixture. Pharmaceutical constituents such as fillers, binders, lubricants, disintegrating agents, surfactants and other pharmaceutically acceptable additives.

Further, the proton pump inhibitor may also be mixed with an alkaline, pharmaceutically acceptable substance (or substances). Such substances can be chosen among, but are not restricted to substances such as the sodium, potassium, calcium, magnesium and aluminium salts of phosphoric acid, carbonic acid, citric acid or other suitable weak inorganic or organic acids; aluminium hydroxide/sodium bicarbonate coprecipitate; substances normally used in antacid preparations such as aluminium, calcium and magnesium hydroxides; magnesium oxide or composite substances, such as  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$ ,  $(\text{Mg}_6\text{Al}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O})$ ,  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$  or similar compounds; organic pH-buffering substances such as trihydroxymethyl-aminomethane, basic amino acids and their salts or other similar, pharmaceutically acceptable pH-buffering substances.

Alternatively, the aforementioned core material can be prepared by using spray drying or spray congealing technique.

#### Enteric coating layer(s)

Before applying the enteric coating layer(s) onto the core material in the form of individual pellets or tablets, the pellets or tablets may optionally be covered with one or more separating layer(s) comprising pharmaceutical excipients optionally including alkaline

compounds such as pH-buffering compounds. This/these separating layer(s), separate(s) the core material from the outer layers being enteric coating layer(s). The separating layer(s) protecting the core material of a proton pump inhibitor should be water soluble or rapidly disintegrating in water.

5

The separating layer(s) can be applied to the core material by coating or layering procedures in suitable equipments such as coating pan, coating granulator or in a fluidized bed apparatus using water and/or organic solvents for the coating process. As an alternative the separating layer(s) can be applied to the core material by using powder coating technique.

10 The materials for the separating layers are pharmaceutically acceptable compounds such as, for instance, sugar, polyethylene glycol, polyvinylpyrrolidone, polyvinyl alcohol, polyvinyl acetate, hydroxypropyl cellulose, methylcellulose, ethyl-cellulose, hydroxypropyl methyl cellulose, carboxymethylcellulose sodium and others, used alone or in mixtures. Additives such as plasticizers, colorants, pigments, fillers anti-tacking and anti-static agents, such as  
15 for instance magnesium stearate, titanium dioxide, talc and other additives may also be included into the separating layer(s).

When the optional separating layer, is applied to the core material it may constitute a variable thickness. The maximum thickness of the separating layer(s) is normally only  
20 limited by processing conditions. The separating layer may serve as a diffusion barrier and may act as a pH-buffering zone. The pH-buffering properties of the separating layer(s) can be further strengthened by introducing into the layer(s) substances chosen from a group of compounds usually used in antacid formulations such as, for instance, magnesium oxide, hydroxide or carbonate, aluminium or calcium hydroxide, carbonate or silicate; composite  
25 aluminium/magnesium compounds such as, for instance  $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$ ,  $(\text{Mg}_6\text{Al}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O})$ ,  $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ , aluminium hydroxide/sodium bicarbonate coprecipitate or similar compounds; or other pharmaceutically acceptable pH-buffering compounds such as, for instance the sodium, potassium, calcium, magnesium and aluminium salts of phosphoric, carbonic, citric or other suitable, weak, inorganic or organic  
30 acids; or suitable organic bases, including basic amino acids and salts thereof. Talc or other

compounds may be added to increase the thickness of the layer(s) and thereby strengthen the diffusion barrier. The optionally applied separating layer(s) is not essential for the invention. However, the separating layer(s) may improve the chemical stability of the active substance and/or the physical properties of the novel multiple unit tableted dosage form.

5

Alternatively, the separating layer may be formed in situ by a reaction between an enteric coating polymer layer applied on the core material an alkaline reacting compound in the core material. Thus, the separating layer formed comprises a salt formed between the enteric coating layer polymer(s) and an alkaline reacting compound which is in the position  
10 to form a salt.

One or more enteric coating layers are applied onto the core material or onto the core material covered with separating layer(s) by using a suitable coating technique. The enteric coating layer material may be dispersed or dissolved in either water or in suitable organic  
15 solvents. As enteric coating layer polymers one or more, separately or in combination, of the following can be used, e.g. solutions or dispersions of methacrylic acid copolymers, cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, polyvinyl acetate phthalate, cellulose acetate trimellitate, carboxymethylethylcellulose, shellac or other suitable enteric coating polymer(s).

20

The enteric coating layers may contain pharmaceutically acceptable plasticizers to obtain the desired mechanical properties, such as flexibility and hardness of the enteric coating layers. Such plasticizers are for instance, but not restricted to triacetin, citric acid esters, phthalic acid esters, dibutyl sebacate, cetyl alcohol, polyethylene glycols, polysorbates or other  
25 plasticizers.

30

The amount of plasticizer is optimized for each enteric coating layer formula, in relation to selected enteric coating layer polymer(s), selected plasticizer(s) and the applied amount of said polymer(s), in such a way that the mechanical properties, i.e. flexibility and hardness of the enteric coating layer(s), for instance exemplified as Vickers hardness, are adjusted so

that the acid resistance of the pellets covered with enteric coating layer(s) does not decrease significantly during compression of pellets into tablets. The amount of plasticizer is usually above 10 % by weight of the enteric coating layer polymer(s), preferably 15 - 50 % and more preferably 20 - 50 %. Additives such as dispersants, colorants, pigments polymers e.g. poly (ethylacrylat, methylmethacrylat), anti-tacking and anti-foaming agents may also be included into the enteric coating layer(s). Other compounds may be added to increase film thickness and to decrease diffusion of acidic gastric juices into the acid susceptible material.

To protect the acid susceptible substance, the proton pump inhibitor, and to obtain an acceptable acid resistance of the dosage form according to the invention, the enteric coating layer(s) constitutes a thickness of approximately at least 10  $\mu\text{m}$ , preferably more than 20  $\mu\text{m}$ . The maximum thickness of the applied enteric coating is normally limited by processing conditions and the desired dissolution profile.

Alternatively the enteric coating layer described above may be used for enteric coating of conventional tablets comprising an acid susceptible proton pump inhibitor. Said enteric coating layered tablet is thereafter presscoated with a granulation comprising the prokinetic compound.

#### Over-coating layer

Pellets covered with enteric coating layer(s) may further be covered with one or more over-coating layer(s). The over-coating layer(s) should be water soluble or rapidly disintegrating in water. The over-coating layer(s) can be applied to the enteric coating layered pellets by coating or layering procedures in suitable equipments such as coating pan, coating granulator or in a fluidized bed apparatus using water and/or organic solvents for the coating or layering process. The materials for over-coating layers are chosen among pharmaceutically acceptable compounds such as, for instance sugar, polyethylene glycol, polyvinylpyrrolidone, polyvinyl alcohol, polyvinyl acetate, hydroxypropyl cellulose, methylcellulose, ethylcellulose, hydroxypropyl methyl cellulose, carboxymethylcellulose

- sodium and others, used alone or in mixtures. Additives such as plasticizers, colorants, pigments, fillers, anti-tacking and anti-static agents, such for instance magnesium stearate, titaniumdioxide, talc and other additives may also be included into the over-coating layer(s). Said over-coating layer may further prevent potential agglomeration of enteric coating
- 5 layered pellets, further protect the enteric coating layer towards cracking during the compaction process and enhance the tableting process. The maximum thickness of the applied over-coating layer(s) is normally limited by processing conditions and the desired dissolution profile.
- 10 The above described over-coating layer may also be used as a tablet filmcoat to obtain tablets of good appearance.

#### Prokinetic preparation

- 15 The active substance(s) in form of one or more prokinetic compound(s) is dry mixed with inactive excipients and the mixture is wet massed with a granulation liquid. The wet mass is dried preferably to a loss on drying of less than 3% by weight. Thereafter the dry mass is milled to a suitable size for the granules, such as smaller than 4 mm, and preferably smaller than 1 mm. Suitable inactive excipients for the prokinetic mixture are for instance lactose,
- 20 corn starch low substituted hydroxypropyl cellulose, microcrystalline cellulose, sodium starch glycolate and crosslinked polyvinyl pyrrolidone. The dry mixture comprising prokinetic compound is wet-mixed with a suitable granulation liquid comprising for instance hydroxy propyl cellulose or polyvinyl pyrrolidone dissolved in purified water or an alcohol or a mixture thereof. Alternatively, the prokinetic agent(s) are dry mixed with
- 25 pharmaceutically acceptable excipients according to above.

- As a further alternative, the prokinetic agent(s) can be applied in a separate layer onto a multiple unit tableted dosage form or surrounding the tablet comprising the proton pump inhibitor. The prokinetic agent(s) is dispersed or dissolved in an aqueous solution optionally
- 30 comprising binders for suspension layering onto the tablet.

### Multiple unit tablets

The enteric coating layered pellets comprising a proton pump inhibitor are mixed with the  
5 granules comprising prokinetic compound and tablet excipients such as fillers, binders, disintegrants, lubricants and other pharmaceutically acceptable additives. The mixture is compressed into a multiple unit tableted dosage form. The compressed tablet is optionally covered with a filmforming agent(s) to obtain a smooth surface of the tablet and further enhance the stability of the tablet during packaging and transport. Such a coating layer may  
10 further comprise additives such as anti-tacking agents, colorants and pigments or other additives to obtain a tablet of good appearance.

Alternatively the enteric coated pellets may be dry mixed with the prokinetic compound and pharmaceutically acceptable tablet excipients according to above, and compressed into  
15 tablets (direct compression).

Suitable lubricants for the tableting process are for instance sodium stearyl fumarate, magnesium stearate and talc.

20 Further, the different active substances may be formulated into different layers, wherein the layer comprising the proton pump inhibitor is in the form of a multiple unit tableted dosage form layered with prepared prokinetic granules. The two layers may be separated by an anti-tacking layer.

25 As a further alternative the proton pump inhibitor is dry mixed with inactive excipients and compressed into a conventional tablet which is coating layered with an enteric coating and optionally a separating layer is applied before the enteric coating. Thereafter the enteric coated tablet is presscoated with a prokinetic preparation. The tablet core may also be formulated as a multiple unit tableted dosage form comprising the proton pump inhibitor,  
30 the tablet is spray coating layered by a suspension comprising the prokinetic agent(s).

The fraction of enteric coating layered pellets constitutes less than 75 % by weight of the total tablet weight and preferably less than 60 %. By increasing the amount of the granules comprising the prokinetic agent the fraction of enteric coating layered pellets of the proton pump inhibitor may be reduced in the multiple unit tableted dosage form. By choosing small enteric coating layered pellets in the formulation according to the present invention, the number of pellets in each tablet can be held high which in turn makes the tablet divisible with retained dosing accuracy.

Thus, the preferred multiple unit tablet formulation consists of enteric coating layered pellets containing one active substance in the form of a proton pump inhibitor, optionally admixed with alkaline reacting compound(s), compressed into tablets together with the prepared prokinetic mixture and optionally tablet excipients. The addition of an alkaline reacting material to the proton pump inhibitor is not necessary, in any sense but such a substance may further enhance the stability of the proton pump inhibitor or some of the alkaline reacting compounds may react in situ with the enteric coating material to form a separating layer. The enteric coating layer(s) is making the pellets of the dosage form insoluble in acidic media, but disintegrating/dissolving in near neutral to alkaline media such as, for instance the liquids present in the proximal part of the small intestine, where dissolution of the proton pump inhibitor is desired. The prokinetic agent(s) may be released in the stomach. The enteric coating layered pellets may further be covered with an overcoating layer before being formulated into the tablet and they may also contain one or more separating layer(s) optionally containing alkaline substance(s).

## Process

The process for the manufacture of the dosage form represents a further aspect of the invention. After formulation of the pellets by spray coating or layering of the proton pump inhibitor onto seeds, or by extrusion/spheronization or granulation, e.g. rotor granulation of homogeneous pellets, the pellets are first optionally covered with the separating layer(s) and

then with the enteric coating layer(s) or a separating layer is spontaneously developed in situ between the alkaline core material and the enteric coating layer material. The coating is carried out as described above and in the accompanying examples. The preparation of the prokinetic mixture is also described above and in the examples. The pharmaceutical  
5 processes can preferably be completely water-based.

The enteric coating layered pellets, with or without an over-coat, are mixed with the prepared prokinetic mixture, optionally tablet excipients and other pharmaceutically acceptable additives and compressed into tablets. Alternatively, the enteric coating layered  
10 pellets may be intimately mixed with tablet excipients and precompressed and further layered with the prepared prokinetic mixture and finally compressed into a tablet. As a further alternative the proton pump inhibitor in form of the active substance may be mixed with tablet excipients and compressed into a tablet which is optionally layered with a separating layer and thereafter enteric coating layered. Said tablet is then presscoated with  
15 the prepared prokinetic mixture. Alternatively, a multiple unit tableted dosage form of the proton pump inhibitor is manufactured as describes above. The multiple unit dosage form is spray coating layered by an aqueous suspension comprising the prokinetic agent(s). The suspension may optionally comprise binders; such as hydroxypropyl methylcellulose, and an alcohol to solve the binder. The proton pump inhibitor in the form of enteric coating layered  
20 pellets may also be filled into a capsule together with the prokinetic substance in the form of a granulation optionally mixed with pharmaceutical excipients.

#### Use of the preparation

25 The dosage forms according to the invention are especially advantageous in the treatment of gastro oesophageal reflux disease and other gastrointestinal disorder. They are administered one to several times a day, preferably once or twice daily. The typical daily dose of the active substances varies and will depend on various factors such as the individual requirements of the patients, the mode of administration and disease. In general each dosage  
30 form will comprise 0.1-200 mg of the proton pump inhibitor and 0.1-100 mg of the



prokinetic compound. Preferably, each dosage form will comprise 10-80 mg of the proton pump inhibitor and 3-80 mg of the prokinetic compound, and more preferably 10-40 mg of proton pump inhibitor and 15 - 40 mg of the prokinetic compound, respectively.

- 5 The multiple unit tablet preparation is also suitable for dispersion in an aqueous liquid with slightly acidic pH-value before being orally administered or fed through a naso-gastric tube.

The invention is illustrated more in detail in the following examples.

10 **Examples**

**Example 1:**

Multiple unit dosage form comprising magnesium omeprazole and mosapride (batch size  
15 500 tablets).

**Core material**

	Magnesium omeprazole	5 kg
	Sugar sphere seeds	10 kg
20	Hydroxypropyl methylcellulose	0.75 kg
	Water purified	20.7 kg

**Separating layer**

	Core material (acc. to above)	10.2 kg
25	Hydroxypropyl cellulose	1.02 kg
	Talc	1.75 kg
	Magnesium stearate	0.146kg
	Water purified	21.4 kg

Enteric coating layer

	Pellets covered with separating layer (acc. to above)	11.9 kg
	Methacrylic acid copolymer (30 % suspension)	19.8 kg
	Triethyl citrate	1.79 kg
5	Mono- and diglycerides (NF)	0.297kg
	Polysorbate 80	0.03 kg
	Water purified	11.64 kg

Over-coating layer

10	Enteric coating layered pellets (acc. to above)	20 kg
	Hydroxypropyl methylcellulose	0.238kg
	Magnesium stearate	0.007kg
	Water purified	6.56 kg

15 Tablets

	Prepared pellets comprising omeprazole (acc. to above)	41.2 g
	Mosapride citrate dihydrate	23.4 g
	Microcrystalline cellulose	138.1 g
	Polyvinyl pyrrolidone crosslinked	2.9 g
20	Sodium stearyl fumarate	0.29 g

Tablet coating solution (for 10 kg tablets)

	Hydroxypropyl methylcellulose	250 g
	Polyethylene glycol 6000	62.5 g
25	Titanium dioxide	62.5 g
	Water purified	2125 g
	Hydrogen pyroxide	0.75 g

Suspension layering was performed in a fluid bed apparatus. Magnesium omeprazole was sprayed onto sugar sphere seeds from a water suspension containing the dissolved binder. The size of sugar sphere seeds were in the range of 0.25 to 0.35 mm.

- 5 The prepared core material was covered with a separating layer in a fluid bed apparatus with a hydroxypropyl cellulose solution containing talc and magnesium stearate. The enteric coating layer consisting of methacrylic acid copolymer, mono- and diglycerides, triethyl citrate and polysorbate was sprayed onto the pellets covered with a separating layer in a fluid bed apparatus. In a fluid bed apparatus enteric coating layered pellets were coated with  
10 a hydroxypropyl methylcellulose solution containing magnesium stearate. The over-coating layered pellets were classified by sieving.

- The enteric coating layered pellets with an over-coating layer, mosapride citrate dihydrate, microcrystalline cellulose, polyvinyl pyrrolidone crosslinked and sodium stearyl fumarate  
15 were dry mixed and compressed into tablets using an excenter tableting machine equipped with 12 mm punches. The amount of omeprazole in each tablet was approx. 10 mg and the amount of mosapride was approx. 30 mg. The tablet hardness was measured to 70-80 N.

The obtained tablets are covered with a conventional tablet filmcoating layer.

20

#### Example 2:

Multiple unit dosage form comprising magnesium omeprazole and mosapride (batch size 500 tablets).

25

#### Mosapride granulation

Mosapride citrate dihydrate	46.8 g
Lactose monohydrate	350 g
Corn starch	184 g
30 Hydroxy propyl cellulose LF	25 g

Water purified	225 g
Hydroxypropyl cellulose (L-HPC)	152 g
Magnesium stearate	7.4 g

5 **Tablets**

Enteric coating layered pellets with an over-coating layer (manufacturing and composition as in example 1)	41.2 g
Mosapride granulation	190 g

10 **Tablet coating solution (for 10 kg tablets)**

Hydroxypropyl methyl cellulose	250 g
Polyethylene glycol 6000	62.5 g
Titaniumdioxid	62.5 g
Water purified	2125 g
15 Hydrogen peroxide	0.75g

Hydroxypropyl cellulose was dissolved in purified water to form the granulation liquid. Mosapride citrate dihydrate, lactose monohydrate and corn starch were dry mixed. The granulation liquid was added to the powder mixture and the mass was wet-mixed. The wet mass was dried in a steam-oven and milled through sieve 1 mm in an oscillating mill equipment. The prepared granulation was mixed with low substituted hydroxypropyl cellulose and magnesium stearate.

The enteric coating layered pellets with an over-coat and prepared granules were mixed and compressed into tablets using an excenter tableting machine equipped with 11 mm punches. The amount of omeprazole in each tablet was approx. 10 mg and the amount of mosapride was approx. 15 mg. Tablet hardness was measured to 30 - 40 N.

The obtained tablets are covered with a conventional tablet filmcoating layer.

Example 3:

Multiple unit dosage form comprising magnesium omeprazole and mosapride (batch size 500 tablets).

5

Core material

	Magnesium omeprazole	10	kg
	Sugar sphere seeds	10	kg
	Hydroxypropyl methylcellulose	1.5	kg
10	Water purified	29.9	kg

Separating layer

	Core material (acc. to above)	20	kg
	Hydroxypropyl cellulose	2	kg
15	Talc	3.43	kg
	Magnesium stearate	0.287	kg
	Water purified	41	kg

Enteric coating layer

20	Pellets covered with separating layer (acc. to above)	24.5	kg
	Methacrylic acid copolymer (30 % suspension)	32.7	kg
	Triethyl citrate	2.94	kg
	Mono- and diglycerides (NF)	0.49	kg
	Polysorbate 80	0.049	kg
25	Water purified	19.19	kg

Over-coating layer

	Enteric coating layered pellets (acc. to above)	37.8	kg
	Hydroxypropyl methylcellulose	0.49	kg
30	Magnesium stearate	0.0245	kg

Water purified 11.6 kg

Tablets

	Prepared pellets comprising omeprazole (acc. to above)	47.45 g
5	Mosapride citrate dihydrate	23.4 g
	Microcrystalline cellulose	163 g
	Polyvinyl pyrrolidone crosslinked	3.3 g
	Sodium stearyl fumarate	0.3 g
10	<u>Tablet coating solution (for 10 kg tablets)</u>	
	Hydroxypropyl methyl cellulose	250 g
	Polyethylene glycol 6000	62.5 g
	Titanium dioxid	62.5 g
	Water purified	2125 g
15	Hydrogen peroxide	0.75 g

The enteric coating layered pellets with an over-coating layer prepared as described in Example 1, mosapride citrate dihydrate, microcrystalline cellulose, polyvinyl pyrrolidone crosslinked and sodium stearyl fumarate were dry mixed and compressed into tablets using an excenter tableting machine equipped with 12 mm punches.

The amount of omeprazole in each tablet was approx. 20 mg and the amount of mosapride was approx. 30 mg. The tablet hardness was measured to 70 N.

The tablets are covered with a conventional tablet filmcoating layer.

Example 4:

Multiple unit dosage form comprising S-omeprazole magnesium salt and mosapride (batch size 300 tablets).

Core material

	S-omeprazole magnesium salt	120	g
	Sugar sphere seeds	150	g
	Hydroxypropyl methylcellulose	18	g
5	Polysorbate 80	2.4	g
	Water purified	562	g

Separating layer

10	Core material (acc. to above)	200	g
	Hydroxypropyl cellulose	30	g
	Talc	51.4	g
	Magnesium stearate	4.3	g
	Water purified	600	g

15

Enteric coating layer

	Pellets covered with separating layer (acc. to above)	250	g
	Methacrylic acid copolymer (30% suspension)	333.7	g
20	Triethyl citrate	30	g
	Mono- and diglycerides (NF)	5	g
	Polysorbate 80	0.5	g
	Water purified	196	g

25 Tablets

	Prepared pellets comprising (s)-omeprazole Mg-salt (acc. to above)	38.2	g
	Mosapride citrate dihydrate	14	g
	Microcrystalline cellulose	98.3	g
	Polyvinyl pyrrolidone crosslinked	2.1	g
30	Sodium stearyl fumarate	0.2	g

Tablet coating solution (for 10 kg tablets)

	Hydroxypropyl methyl cellulose	250	g
	Polyethylene glycol 6000	62.5	g
5	Titaniumdioxid	62.5	g
	Water purified	2125	g
	Hydrogen peroxide	0.75	g

10 Suspension layering was performed in a fluid bed apparatus. S-Omeprazole magnesium salt was sprayed onto sugar sphere seeds from a water suspension containing the dissolved binder and polysorbate 80. The size of sugar sphere seeds were in the range of 0.25 to 0.35 mm.

15 The prepared core material was covered with a separating layer in a fluid bed apparatus with hydroxypropyl cellulose solution containing talc and magnesium stearate. The enteric coating layer consisting of methacrylic acid copolymer, mono-and diglycerides, triethyl citrate and polysorbate was sprayed onto the pellets covered with a separating layer in a fluid bed apparatus. The enteric coating layered pellets were classified by sieving.

20 The enteric coating layered pellets, mosapride citrate dihydrate, microcrystalline cellulose, polyvinyl pyrrolidone crosslinked and sodium stearyl fumarate were mixed and compressed into tablets using an excenter tableting machine equipped with 12mm punches. The amount of S-omeprazole in each tablet was approx. 20 mg and the amount of mosapride was approx. 30 mg. The tablet hardness was measured to 65 N.

25

The tablets are covered with a conventional tablet filmcoating layer.



"Acid resistance" i.e. % left after exposure to 0.1 N HCl for 2 hrs	
	Tablets
Ex 1	97%
Ex 2	90%
Ex 3	102%
Ex 4	104%

Example 5:

5

Multiple unit dosage form comprising lanzoprazole and mosapride (batch size 500 tablets).

Core material

	Lanzoprazole	400 g
10	Sugar sphere seeds	400 g
	Hydroxypropyl methylcellulose	80 g
	Sodium laurylsulfate	3 g
	Water purified	1500 g

15 Separating layer

	Core material (acc. to above)	400 g
	Hydroxypropyl cellulose	40 g
	Talc	69 g
	Magnesium stearate	6 g
20	Water purified	800 g

Enteric coating layer

	Pellets covered with a separating layer (acc. to above)	400 g
	Methacrylic acid copolymer (30 % suspension)	667 g
	Triethyl citrate	60 g
5	Mono- and diglycerides (NF)	10 g
	Polysorbate 80	1 g
	Water purified	420 g

Tablets

10	Prepared pellets comprising lansoprazole (acc. to above)	47 g
	Mosapride citrate dihydrate	46.8 g
	Microcrystalline cellulose	261 g
	Polyvinyl pyrrolidone crosslinked	5 g
	Sodium stearyl fumarate	0.5 g

15

Tablet coating solution (for 10 kg tablets)

	Hydroxypropyl methylcellulose	250 g
	Polyethylene glycol 6000	62.5 g
	Titanium dioxid	62.5 g
20	Water purified	2125 g
	Hydrogen peroxide	0.75 g

Suspension layering was performed in a fluid bed apparatus. Lansoprazole was sprayed onto the sugar sphere seeds from a suspension containing the dissolved binder in a water solution. Pellets covered with separating layer and enteric coating layer were produced as in example 1.

The enteric coating layered pellets, mosapride citrate dihydrate, microcrystalline cellulose, polyvinyl pyrrolidone crosslinked and sodium stearyl fumarate were dry mixed and compressed into tablets using an excenter tableting machine equipped with 10 mm punches.

The amount of lansoprazole in each tablet was approx. 10 mg and the amount of mosapride was approx. 30 mg. The tablet hardness was measured to 70 N.

The tablets are covered with a conventional tablet filmcoating layer.

5

Example 6:

Magnesium omeprazole and mosapride presscoated tablets (batch size 10.000 tablets).

10 Omeprazole tablets

Mg-omeprazole	112.5 g
Mannitol	287 g
Microcrystalline cellulose	94 g
Sodium starch glycolate	30 g
15 Hydroxypropyl methylcellulose	30 g
Talc	25 g
Microcrystalline cellulose	31 g
Sodium stearyl fumarate	12.5 g
Water purified	200 g

20

Solution for separating layer (for 10 kg tablets)

Hydroxypropyl methylcellulose	300 g
Hydrogen peroxide (30%)	0.003 g
Water purified	2700 g

25

Solution for enteric coating layer (for 10 kg tablets)

Methacrylic acid copolymer dispersion (30%)	2450 g
Polyethylene glycol 400	80 g
Titanium dioxide Colour	100 g
30 Water purified	1960 g

Presscoated tablet

Mg-Omeprazole tablets 10.000 tabl

Mosapride granulation

5 (manufacturing and composition as in example 2) 3800 g

Tablet coating solution (for 10 kg tablets)

Hydroxypropyl methylcellulose 250 g

Polyethylene glycol 6000 62.5 g

10 Titaniumdioxid 62.5 g

Water purified 2125 g

Hydrogen peroxide 0.75 g

15 Magnesium omeprazole, mannitol, microcrystalline cellulose, sodium starch glycolate and hydroxypropyl methyl cellulose are dry mixed. The powder mixture is moistened with water purified. The granulation is dried and milled through sieve 1 mm in a suitable mill. The prepared granules comprising proton pump inhibitor is mixed with talc, microcrystalline cellulose and sodium stearyl fumarate and compressed into tablets using a rotary tableting machine equipped with 5 mm punches.

20

The obtained tablets are coated layered with a separating layer and an enteric coating layer. Said tablets are then presscoated with mosapride granulation using a presscoating machine equipped with 11 mm punched.

25 The tablets are covered with a conventional tablet filmcoating layer.

Example 7:

A capsule formulation comprising magnesium omeprazole and mosapride (batch size 100 capsules).

5

Capsules

Enteric coating layered pellets with an over-coating layer 9.49 g  
(manufacturing and composition as in example 3)

Mosapride granulation 38 g

10 (manufacturing and composition as in example 2)

Enteric coating layered pellets and mosapride granulation are filled into capsules, size 00.  
The amount of omeprazole in each capsule is approx. 20 mg and the amount of mosapride  
is approx. 15 mg.

15

Example 8:

Multiple unit dosage form comprising magnesium omeprazole with a tablet coating layer  
comprising mosapride (batch size 1 000 tablets).

20

Tablets

Enteric coating layered pellets with an overcoat 82.4 g  
(manufacturing and composition as in example 1)

Microcrystalline cellulose 179.2 g

25 Polyvinyl pyrrolidone crosslinked 3.7 g

Sodium stearyl fumarate 0.4 g

Mosapride coating layer suspension

Mosapride citrate dihydrate 23.4 g

30 Hydroxypropyl methyl cellulose 13.4 g

Ethanol 99 %	132.5 g
Water purified	132.5 g

Tablet coating solution (for 10 kg tablets)

5 Hydroxypropyl methylcellulose	250 g
Polyethylene glycol 6000	62.5 g
Titanium dioxid	62.5 g
Water purified	2125 g
Hydrogen peroxide	0.75g

10

The enteric coating layered pellets are mixed with microcrystalline cellulose, polyvidone and sodium stearyl fumarate and compressed into tablets using an excenter tableting machine equipped with 9 mm punches. The tablets are then coated layered in a fluid bed apparatus with the suspension comprising mosapride. The amount of omeprazole in each tablet is

15 approx. 10 mg and the amount of mosapride is approx. 15 mg.

Finally the tablets are covered with a conventional tablet filmcoating layer.

The best mode to practise the invention is according to compositions described in Examples

20 1 and 4.

The enteric coating layered pellets comprising a proton pump inhibitor may also be prepared as described in the following examples.

25 Example 9:

Preparation of enteric coating layered pellets by extrusion/spheronization.

Core material

	Magnesium omeprazole	600 g
	Mannitol	1000 g
	Microcrystalline cellulose	300 g
5	Hydroxypropyl cellulose	100 g
	Sodium lauryl sulphate	6 g
	Water purified	802 g

Separating layer

10	Core material	400 g
	Hydroxypropyl methylcellulose	48 g
	Water purified	960 g

Enteric coating layer

15	Pellets covered with separating layer	200 g
	Methacrylic acid copolymer	100 g
	Triethyl citrate	30 g
	Mono- and diglycerides (NF)	5 g
	Polysorbate 80	0.5 g
20	Water purified	309 g

Sodium lauryl sulphate is dissolved in purified water to form the granulation liquid.

Magnesium omeprazole, mannitol, microcrystalline cellulose and hydroxypropyl cellulose are dry-mixed. The granulation liquid is added to the powder mixture and the mass is wet-  
25 mixed.

The wet mass is forced through an extruder equipped with screens of size 0.5 mm. The extrudate is spheronized on a friction plate in a spheronizing apparatus. The core material is dried in a fluid bed dryer and classified. The prepared core material is covered by a

separating layer in a fluid bed apparatus with a hydroxypropyl methylcellulose/water solution.

The enteric coating layer is applied to the pellets covered with separating layer from an aqueous dispersion of methacrylic acid copolymer plasticized with triethyl citrate to which a mono- and diglycerides/polysorbate dispersion has been added. The pellets are dried in a fluid bed apparatus.

Example 10:

10

Preparation of enteric coating layered pellets by powder.

Core material

	Magnesium omeprazole	1 500 g
15	Sugar sphere seeds	1 500 g
	Hydroxypropyl methylcellulose	420 g
	Aerosil®	8 g
	Water purified	4 230 g

20 Separating layer

	Core material	500 g
	Hydroxypropyl cellulose	40 g
	Talc	67 g
	Magnesium stearate	6 g
25	Water purified	800 g

Enteric coating layer

	Pellets covered with separating layer	500 g
	Methacrylic acid copolymer	200 g



Triethyl citrate	60 g
Water purified	392 g

5 Magnesium omeprazole, part of the hydroxypropyl methylcellulose and Aerosil® are dry-mixed forming a powder. Sugar sphere seeds (0.25-0.40 mm) are layered with the powder in a centrifugal fluidized coating granulator while spraying a hydroxypropyl methylcellulose solution (6 %, w/w).

10 The prepared core material is dried and covered by a separating layer in a centrifugal fluidized coating-granulator. A fluid bed apparatus is used for enteric coating layering.

Example 11:

Preparation of enteric coating layered pellets with of silicon dioxide seeds.

15

Core material

Magnesium omeprazole	8.00 kg
Silicon dioxide	8.00 kg
Hydroxypropyl methylcellulose	1.41 kg
20 Sodium lauryl sulphate	0.08 kg
Water purified	28.00 kg

Separating layer

Core material	10.00 kg
25 Hydroxypropyl methylcellulose	0.80 kg
Water purified	10.00 kg

Enteric coating layer

Pellets covered with separating layer	300 g
30 Methacrylic acid copolymer	124 g

Polyethylene glycol 400	25 g
Mono- and diglycerides (NF)	3 g
Polysorbate 80	1 g
Water purified	463 g

5

Suspension layering is performed in a fluid bed apparatus. Magnesium omeprazole is sprayed onto the silicon dioxide seeds from a water suspension containing the dissolved binder and a surface active ingredient.

- 10 The prepared core material is covered with a separating layer in a fluid bed apparatus with a hydroxypropyl methylcellulose solution. The enteric coating layer consisting of methacrylic acid copolymer, mono- and diglycerides, polyethylene glycol 400 and polysorbate is sprayed onto the pellets covered with separating layer in a fluid bed apparatus.

15 Example 12:

Preparation of enteric coating layered pellets.

Enteric coating layer

- 20 Pellets covered with separating layer  
(manufacturing and composition

as in example 10)	500 g
Methacrylic acid copolymer	250 g
Polyethylene glycol 6000	75 g
25 Mono- and diglycerides (NF)	12.5 g
Polysorbate 80	1.2 g
Water purified	490 g

Example 13:

Preparation of enteric coating layered pellets.

5 Enteric coating

Pellets covered with separating layer 500 g

(manufacturing and composition as in example 1)

Hydroxypropyl methylcellulose phthalate 250 g

Cetanol 50 g

10 Ethanol (95%) 1000 g

Acetone 2500 g

Example 14:

15 Preparation of enteric coating layered pellets.

Core material

Omeprazole 225 g

Mannitol 1425 g

20 Hydroxypropyl cellulose 60 g

Microcrystalline cellulose 40 g

Lactose anhydrous 80 g

Sodium lauryl sulphate 5 g

Disodium hydrogen phosphate dihydrate 8 g

25 Water purified 350 g

Separating layer

Core material 300 g

Hydroxypropyl cellulose 30 g

30 Talc 51 g

Magnesium stearate 4 g

Enteric coating layer

	Pellets covered with separating layer	300 g
5	Methacrylic acid copolymer	140 g
	Triethyl citrate	42 g
	Mono- and diglycerides (NF)	7 g
	Polysorbate 80	0.7 g

- 10 The dry ingredients for producing the core material are well mixed in a mixer. Addition of granulation liquid is made and the mixture is kneaded and granulated to a proper consistency. The wet mass is pressed through an extruder screen and the granules are converted into a spherical form in a spheronizer. The core material is dried in a fluid bed apparatus and classified into a suitable particle size range, e.g. 0.5 - 1.0 mm. The prepared
- 15 core material is covered with a separating layer and enteric coating layered as described in previous examples.

Preparation of active substance

- 20 Magnesium omeprazole used in some of the examples is produced according to the process described in WO95/01977, the single enantiomers of omeprazole salts are prepared as described in WO94/27988 and omeprazole is produced according to the process disclosed in EP-A1 0005129. These documents are hereby incorporated in a whole by reference.

CLAIMS

1. An oral pharmaceutical dosage form comprising an acid susceptible proton pump inhibitor together with at least one prokinetic agent and optionally pharmaceutically acceptable excipients, characterized in that the dosage form is in the form of a fixed unit dosage form comprising at least two pharmaceutically active components, and wherein the proton pump inhibitor is protected by an enteric coating layer.
2. A dosage form according to claim 1, wherein the dosage form is a tablet formulation.
3. A dosage form according to claim 1, wherein the dosage form is a capsule formulation.
4. A dosage form according to claim 1, wherein the proton pump inhibitor is protected by two layers, an enteric coating layer and a layer separating the enteric coating from the proton pump inhibitor.
5. A dosage form according to claim 1, wherein the dosage form comprises a proton pump inhibitor and one prokinetic agent.
6. A dosage form according to claim 1, wherein the proton pump inhibitor is omeprazole, one of its single enantiomer or an alkaline salt thereof.
7. A dosage form according to claim 6, wherein the proton pump inhibitor is S-omeprazole magnesium salt.
8. A dosage form according to claim 1, wherein the proton pump inhibitor is lansoprazole, one of its single enantiomer or an alkaline salt thereof.

9. A dosage form according to one of claims 6 - 8, wherein the prokinetic agent is mosapride.
10. A dosage form according to one of claims 6 - 8, wherein the prokinetic agent is cisapride.
11. A dosage form according to claim 1, wherein the amount of proton pump inhibitor is in the range of 10-80 mg and the amount of prokinetic agent(s) is in the range of 3-80 mg.
12. A dosage form according to claim 1, wherein the amount of proton pump inhibitor is in the range of 10-40 mg and the amount of prokinetic agent(s) is in the range of 15-40 mg.
13. A tableted dosage form according to claim 2, wherein the dosage form consists of two separate layers, one layer comprising a proton pump inhibitor and the other layer comprising one or more prokinetic agents.
14. A tableted dosage form according to claim 2, wherein the tablet formulation is a multiple unit tableted dosage form comprising the proton pump inhibitor in the form of enteric coating layered pellets compressed together with a prokinetic preparation into a tablet, whereby the enteric coating layer covering the pellets has mechanical properties such that the tableting of the pellets together with the prokinetic granulation and optionally pharmaceutically acceptable excipients does not significantly affect the acid resistance of the enteric coating layered pellets.
15. A tableted dosage form according to claim 14, wherein the acid resistance of the enteric coating layered pellets is in coherence with the requirements on enteric coating layered articles defined in the United States Pharmacopeia.

16. A tableted dosage form according to 14, wherein the acid resistance of the enteric coating layered pellets does not decrease more than 10 % during the compression of the pellets into the multiple unit tableted dosage form.
- 5 17. A tableted dosage form according to claim 14, wherein the enteric coating of the pellets comprises a plasticized enteric coating layer material.
18. A tableted dosage form according to claim 14, wherein the enteric coating layered pellets are further covered with an over-coating layer comprising pharmaceutically  
10 acceptable excipients.
19. A tableted dosage form according to claim 14, wherein the tablet is divisible.
20. A tableted dosage form according to claim 19, wherein the tablet is dispersible to a  
15 slightly acidic aqueous suspension comprising a prokinetic agent and enteric coating pellets of a proton pump inhibitor.
21. A tableted dosage form according to claim 2, wherein the tablet is an enteric coating layered tablet comprising the proton pump inhibitor surrounded by a layer comprising the  
20 prokinetic agent.
22. A tableted dosage form according to claim 14, wherein a multiple unit tableted dosage form comprising the proton pump inhibitor is layered with a separate layer comprising the prokinetic agent, or the multiple unit tableted dosage form is surrounded by  
25 a layer comprising the prokinetic agent.
23. A process for the manufacture of a fixed dosage form comprising a proton pump inhibitor and one or more prokinetic agents in a capsule, characterized in that the proton pump inhibitor is prepared in the form of enteric coating layered pellets and the pellets are

filled into a capsule together with the prokinetic agent(s) optionally mixed with pharmaceutically acceptable excipients.

24. A process for the manufacture of a fixed dosage form comprising a proton pump inhibitor and one or more prokinetic agents in a multiple unit tableted dosage form, characterized in that the proton pump inhibitor is prepared in the form of enteric coating layered pellets and these pellets are mixed with prepared prokinetic mixture and optionally pharmaceutically acceptable tablets excipients whereafter the mixture is compressed into a multiple unit tablet without giving any significant change of the acid resistance of the enteric coating layer.

25. A process for the manufacture of a fixed dosage form comprising a proton pump inhibitor and one or more prokinetic agent(s) in an enteric coating layered tablet characterized in that the proton pump inhibitor is admixed with tablet excipients and pre-compressed into a tablet, whereafter tablet is covered with an enteric coating layer and that optionally a separating layer is applied before the enteric coating layer, and the prokinetic agent(s) mixed with pharmaceutically acceptable excipients are thereafter presscoated onto the enteric coating layered tablet.

26. A process for the manufacture of a fixed dosage form comprising a proton pump inhibitor and one or more prokinetic agents in a multiple unit tableted dosage form, characterized in that the proton pump inhibitor is prepared in the form of enteric coating layered pellets and these pellets are mixed with pharmaceutically acceptable tablet excipients and the dry mixture is compressed into a multiple unit tablet without giving any significant change of the acid resistance of the enteric coating layer and whereafter the multiple unit tableted dosage form is spray coating layered by an aqueous suspension of the prokinetic agent(s), or the multiple unit tableted dosage form is layered with a separate layer comprising the prokinetic agent(s) in admixture with pharmaceutically acceptable excipients.



27. A method for the treatment of disorders associated with gastro oesophageal reflux diseases in mammals and man by administering to a host in need thereof a therapeutically effective dose of a multiple unit tableted dosage form according to any of claims 1 to 22.

5 28. A method according to claim 27, wherein the disorder is a gastric disorder associated with gastro oesophageal reflux diseases.

29. Use of a dosage form according to any of claims 1 to 22 for the manufacture of a medicament for treating disorders associated with gastro oesophageal reflux diseases.

10

30. Use according to claim 29 wherein the disorder is a gastric disorder associated with gastro oesophageal reflux diseases.

Fig. 1




Fig. 2




Fig. 3





Fig. 4



# 1

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 96/01736

### A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A61K 45/06, A61K 31/44, A61K 31/445, A61K 9/20, A61K 9/26, A61K 9/48  
According to International Patent Classification (IPC) or to both national classification and IPC

### B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAPLUS, WPI, WPIL, CLAIMS, USFULLTEXT, EMBASE

### C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THE NEW ENGLAND JOURNAL OF MEDICINE, October 1995, Alberto Pilotto, M.D. et al: "A Comparison of five maintenance therapies for reflux esophagitis", page 1106, abstract; page 1109, col. 2, line 11-21, line 27-39 --	1-31
A	WO 9501803 A1 (MERCK & CO., INC.), 19 January 1995 (19.01.95), page 2, line 5 - line 29; page 10, line 17 - page 11, line 7 --	1-31
A	EP 0247983 A2 (AKTIEBOLAGET HÄSSLE), 2 December 1987 (02.12.87), page 4, line 25 - line 2; page 8, line 22 - line 32 --	13-31

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

- \* Special categories of cited documents
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

9 April 1997

22 -04- 1997

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Anneli Jönsson  
Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 96/01736

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0365947 A1 (PHARMACIA AB), 2 May 1990 (02.05.90), page 3, line 41 - line 46; page 4, line 42 - line 57  -- -----	13-31

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 96/01736

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 27-28  
because they relate to subject matter not required to be searched by this Authority, namely:  
  

**Remark: Claims 27-28 are directed to methods of treatment of the human or animal body by surgery or by therapy/diagnostic methods practised on the human or animal body/Rule 39.1(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions**
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

04/03/97

International application No.

PCT/SE 96/01736

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9501803 A1	19/01/95	AU 7397194 A EP 0707492 A JP 8512322 T	06/02/95 24/04/96 24/12/96
EP 0247983 A2	02/12/87	SE 0247983 T3 AR 240250 A AT 140387 T AU 601974 B AU 7191287 A CA 1292693 A CY 1810 A DE 3751860 D,T DE 3783394 A DK 169988 B EP 0496437 A,B SE 0496437 T3 EP 0567201 A ES 2006457 T ES 2091971 T GB 2189698 A HK 135294 A HR 920854 A IE 61416 B JP 1863556 C JP 5294831 A JP 62258320 A LT 1683 A LT 3699 B LV 10357 B NO 174239 B,C SG 154294 A SI 8710681 A SU 1820837 A US 4786505 A	30/03/90 15/08/96 27/09/90 05/11/87 03/12/91 20/10/95 21/11/96 18/02/93 24/04/95 29/07/92 27/10/93 01/01/94 16/11/96 04/11/87 09/12/94 31/10/94 02/11/94 08/08/94 09/11/93 10/11/87 25/07/95 26/02/96 20/04/96 27/12/93 17/03/95 31/10/96 07/06/93 22/11/88
EP 0365947 A1	02/05/90	SE 0365947 T3 AU 612525 B AU 4365089 A CA 2000932 A DE 68907177 T ES 2055775 T HK 123394 A IE 62640 B JP 2164821 A LV 10382 B NO 179478 B,C PT 92103 B SE 8803822 A SG 123894 A US 5178868 A	11/07/91 03/05/90 26/04/90 13/01/94 01/09/94 18/11/94 22/02/95 25/06/90 20/12/95 08/07/96 09/08/95 26/10/88 17/03/95 12/01/93

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**